

Fate of some heavy metals in sandy soil amended with sewage sludge and their accumulation in plants

Badawy, S.H¹. and El-Motaium², R.A.

1-Soil Science Department, Faculty of Agriculture., Cairo University., Giza, Egypt.
Email:<u>shbadawy@main-scc.cairo.eun.eg</u>
2- Plant Research Dept. Nuclear Research Center, Atomic Energy Authority Abou Zaable, Cairo, Egypt.
E-mail: rawia@dns.claes.sci.eg

Abstract

A pot experiment was conducted using tomato plants cultivated in sandy soil amended with different rates (0, 20, 40, 60, 80 t/ha) of sewage sludge collected from El-Gabal El-Asfar farm, Cairo. The same treatments were repeated for non-cultivated soil. Soil samples were collected at different periods (0, 3, 6, 9 and 12 months) to evaluate the release and accumulation of Cu, Zn, Cd, Pb in soil and plant. The results demonstrate that soil pH and CaCO₃ content decreased with increasing sewage sludge application rate and time. Whereas, soil organic matter increased with increasing the application rate and decreased with time. The highest decrease in soil pH and CaCO₃ was found after six months from the beginning of the experiment for both cultivated and non-cultivated soils. However, the decrease in cultivated soil was more than the non-cultivated one. The total content of Cu, Zn, Cd and Pb in soil increased in response to the increase in sludge application rate. No significant changes were observed in soil total metals in relation to the sampling time. The distribution of the different chemical fractions of Cu, Zn, Cd, Pb of the soil are as follows: residual> organic> oxides> carbonates> exchangeable. The mobile fractions and mobile factors (MF) increased up to 6 months from the beginning of the experiment, then decreased. The concentrations of DTPAextractable metals (Cu, Zn, Cd, Pb) in soil increased with the increase in sludge application rate. The values of DTPA-extractable metals Cu, Zn, Cd and Pb were higher in cultivated than in noncultivated soil until 6 months, then they decreased. The highest concentrations of extractable metals were found after 6 months from the beginning of the experiment. The concentrations of Cu, Zn, Cd, Pb in tomato leaves and fruits increased with the increase in sewage sludge application rate. These metals were higher in leaves than in fruits.

Keywords: Heavy Metals, Mobility, Fractionations, Accumulation, Distribution

Introduction

Municipal and industrial sludges can have a beneficial effect on plant growth and crop yields when applied to soil because of their organic matter and nutrients content. However, sludges may also have concentrations of trace elements (e.g., Cd, Cr, Ni, Cu, Pb, and Zn) that are of environmental concern. Evaluation of the potential mobility and biological uptake of trace metal contamination in soil requires chemical or biological methods to quantify the fraction of soil contaminants available for biological uptake. Such information is important for predictive models and risk assessments in systems where the soil-plant pathway is a key contributor to potentially harmful effect to plants and animals. Single and sequential extraction methods have been used to estimate the pollutant concentrations available for plants uptake (He and Singh, 1993; Narwal and Singh, 1998; Singh et al., 1995, 1998, Badawy and Helal, 1997). Several extractants have been used to assess the potential bioavailability of heavy metals for plant uptake (El-Sokkary, 1979; Christensen and Huang, 1999; Kennedy et al., 1997; Aboulroos et al., 1996). According to Pickering (1998), multistep extraction procedures can obtain more detailed information about the status of trace

metals in soils relative to single extraction methods. Furthermore, the early stages to single extraction sequence obtain information comparable to that obtained from single extraction methods. The fraction of metals that is readily displaced from the reversible sorption phases can be regarded as mobile fractions. These fractions can be distinguished from those being irreversible sorbed and released after dissolution in the later stages in the extraction scheme (Salbu et al., 1998). Little work was done to study the effect of time on the distribution of heavy metals among the extractable fractions (Sposito et al., 1983; Aboulroos et al., 1991). The aim of the present work is to evaluate the effect of sewage sludge application and the time factor on: i) the changes of the soil general characteristics. ii) the distribution of the various fractions of Cu, Zn, Cd, and Pb in the soil. iii) the accumulation of these metals in tomato leaves and fruits.

Materials and Methods

A pot experiment was conducted at the plant science department experimental farm, NRC, Anshas. Sandy soil (2.7% clay and 13% WHC) was used and amended with different application rates of sewage sludge collected from El-Gabal El-Asfar farm, Cairo-Egypt (the analyses of sewage sludge were; pH, 6.64; EC, 4.27; CaCO₃, 3.35%; OM, 44%; total Zn, Cu, Cd, and Pb were 760, 219, 5.2, 278 mg/kg, respectively). Treatments were arranged in a Randomized Complete Block Design. Each block included ten treatments (five cultivated with tomato plants (GS cultivar) as a biological indicator for heavy metals accumulation, and five un-cultivated) including a control treatment (recommended chemical fertilizer was applied at a rate of 286 Kg N/ha as ammonium sulphate, 143 Kg P_2O_5 /ha as super phosphate and 167 Kg K₂O/ha as potassium sulphate). Each treatment was replicated four times. Tomato seedlings, GS cultivar, were purchased from a local nursery. Each pot was filled with 5 Kg soil. Distilled water was used for irrigation throughout the growing period and the moisture conditions were kept at 65 to 70 % of water-holding capacity (WHC) of the soil. The experiment was repeated for another year.

Soil Sampling and Analysis:

Soil samples were collected at different periods from the starting time (0, 3, 6, 9, and 12 months). The samples were air dried, crushed to pass through a 2-mm sieve, and stored for analysis. The general characteristics of soils and sewage sludge; pH, electrical conductivity (EC), organic matter (OM), water holding capacity (WHC), calcium carbonate content (CaCO₃), and clay content were determined using the standard methods outlined by Dewis and Freitas (1970) and Jackson (1967).

Total Metal Content:

One-gram (dry weight) soil samples were digested using 12 ml of a mixture of $HClO_4$ and HF (5:1 v/v) in platinum crucible (Jackson, 1967). The crucible was brought to near dryness, before adding a further 12 ml of the acid mixture, and again burning to near dryness. Finally, 1ml of $HClO_4$ was added and the sample was evaporated until the appearance of white fumes. The residue was then dissolved in 5 ml of concentrated HCl before making up to 25 ml with deionised water. Copper, Zinc, and lead were analyzied by Flame Atomic Absorption spectrophotometer and Cadmium by Graphite Furnace Atomic Absorption Spectrophotometer (GF-AAS).

Sequential Chemical Extraction:

The sequential extraction procedure of Tessier et. al.(1979) was used to partition Cu, Zn,Cd, and Pb in soils into five operationally defined fractions; 1) The exchangeable fraction: soils were extracted with 1M NaOAc, pH 8.2 at room temperature for one hour with continuous agitation; 2) The carbonate bound fraction: the reside from 1) was leached at room temperature for five hrs

continuous agitation with 1M NaOAc, pH 5.0 ; 3) The Fe and Mn oxides bond fraction: The residue from 2) was extracted with 0.04 M NH₂ OH.HCl in 25% (v/v) HOAc at 96 \pm 3 °C with occasional agitation for 5 hrs.; 4) The organic matter bound fraction: The residue from 3) was extracted with 0.02 M HNO₃ and 30% H₂O₂, pH 2.0 , for 2 hrs with occasional agitation at 85 \pm 2 °C ; then 30% H₂O₂ for 3 hrs with intermittent agitation at 85 \pm 2 °C ; after cooling 3.2 M NH₄OAc in 20% (v/v) HNO₃ was added.; 5) The residue fraction: the residue from 4) was digested with HF- HClO₄ using the same procedure employed for total. The resulting supernatants from each fractionation step were filtered before analysing the filtrates for metals by flame atomic absorption spectrophotometer and GF-ASS for Cd. using the standard addition technique.

DTPA-Extractable metal:

'Available' Cd, Co, Ni, and Pb were determined for all samples using 0.005 M DTPA (diethylene triamine penta acetic acid) using the methods of Lindsay and Norvell (1978).

Plant Sampling and Analysis:

Tomato plants from each pot, were collected at maturity stage (at early bloom when vegetative growth was completed according to Reisenauer, 1978), then leaves were separated. They were washed with tap water, rinsed with deionized water, oven dried at 60 °C, the dry weight was recorded. The leaves were mixed to form a composite sample, then ground using a micro-mill grinder. The dried samples were stored for metal analyses. Tomato fruits were collected, as they ripe. Representative fruit samples were oven dried at 60 °C for analysis. Heavy metals (Cu, Zn, Cd, and Pb) content of plant materials were measured after digestion with HNO3 - H2SO4 - HClO4 acids mixture (Jackson, 1967). Copper, Zn, and Pb concentrations were measured using the Atomic Absorption Spectrophotometer and the GF-ASS for Cd. using the standard addition technique.

Statistical Analysis:

The data represent the results of two years, were subjected to analysis of variance, LSD, and correlation coefficient (r) using the microcomputer statistical analysis package (MicroStat).

Results and Discussion

General properties of the soil:

Data of Table 1 show that the increases in sewage sludge application rate have resulted in a continuous increase in organic matter content (OM), and electric conductivity (EC), and partly decreased soil pH and CaCO₃ content; the reduction in soil pH, approximately by one unit, CaCO₃ content by 25% and the increase in OM content from 5 to 7 fold, and in EC values by 2 times, comparing the control with the highest sewage sludge application rate at 0 time. These results agree with Aboulroos et. al., 1989; Topper and Sabey, 1986. A glance at Fig. 1. shows that time and vegetation factors affected the changes of OM, EC, pH, and CaCO₃ of the soil. However, the changes were continuously decreased by increased time, the maximum reduction was after 6 months in cultivated soils but, soil organic matter was reduced more at un-cultivated soil. The reduction was approximately 0.7 and 0.6 unit for soil pH, 60 and 30 % for soil CaCO3, and 33 and 40 % for soil OM in cultivated and uncultivated soils, respectively. This could be due to the release of organic acids from sewage sludge and root exudates (El-Motaium and Badawy, 2000).

Soil characteristics		Time (month)									
		0	3	6	9	12	0	3	6	9	12
Treatme	ent			pН			CaCO ₃ (%)				
	Control	8.50	8.40	8.32	8.25	8.28	2.19	2.05	1.79	1.65	1.62
	20t/ha	8.15	8.02	7.65	7.78	7.89	1.61	1.56	1.12	1.23	1.24
Cultivated	40t/ha	7.95	7.54	7.19	7.28	7.25	2.21	1.95	1.85	1.76	1.68
	60t/ha	7.80	7.46	7.25	7.37	7.35	1.77	1.44	1.15	0.79	0.76
	80t/ha	7.65	7.11	7.00	7.16	7.20	1.65	1.45	0.85	0.68	0.51
	Control	8.54	8.44	8.45	8.40	8.40	2.32	2.25	2.10	2.10	2.00
Uncultivated	20t/ha	8.26	8.15	7.85	7.83	7.93	1.42	1.38	1.27	1.23	1.24
	40t/ha	8.00	7.50	7.23	7.35	7.40	1.60	1.60	1.38	1.59	1.55
	60t/ha	7.90	7.41	7.35	7.30	7.46	2.05	1.63	1.45	1.10	0.95
	80t/ha	7.80	7.26	7.22	7.41	7.44	1.65	1.50	1.10	0.80	0.65
		OM (%)					EC (mmohs/cm)				
	Control	0.41	0.45	0.56	0.56	0.54	0.66	0.61	0.95	0.65	0.69
	20t/ha	0.92	0.85	0.74	0.69	0.65	0.82	0.61	0.95	0.66	0.54
Cultivated	40t/ha	1.32	1.10	1.00	1.00	0.87	1.20	1.00	1.10	0.79	0.6
	60t/ha	1.86	1.45	1.25	1.20	1.00	1.7	1.31	1.18	1.35	1.19
	80t/ha	3.20	2.67	2.15	2.00	1.58	1.84	1.50	1.44	1.12	1.02
	Control	0.42	0.41	0.42	0.40	0.39	0.69	0.68	1.00	0.63	0.64
Uncultivated	20t/ha	0.84	0.65	0.50	0.41	0.40	0.80	0.78	1.06	0.63	0.64
	40t/ha	1.23	1.00	0.80	0.77	0.70	1.11	1.09	1.40	0.73	0.74
	60t/ha	1.57	1.35	0.98	0.85	0.72	1.57	1.54	1.25	1.67	1.68
	80t/ha	2.30	2.10	1.25	1.16	1.00	2.01	1.97	1.78	1.37	1.38

Table (1): changes in soil characteristics with increasing the rate of sewage sludge application and time.

Distribution of heavy metals in soil:

The total contents of Cu, Zn, Cd, and Pb in soil are given in Table (2). The results show that the levels of heavy metals in soil increased with increasing application rate of sewage sludge. Comparing the content of heavy metals of the present study with maximum permitted metal loadings in the soils, established by the USEPA-503 regulation (McBride, 1995) showed that the permitted levels of Cu (750), Zn (1400), Cd (20), and Pb (150 mg/kg) were 30, 10, 15, and 5 times of their levels in the present soil at highest application of sewage sludge.

Table (2): changes of the total content of various heavy metals (mg/kg) in sewage sludge treated soil.

biuuz	50 filoatea boll .			
Treatment	Cu	Zn	Cd	Pb
Control	$7.80 \pm 0.65*$	61.90 ± 3.73	0.45 ± 0.031	4.10 ± 0.25
20t/ha	16.1 ± 1.27	66.60 ± 0.96	0.60 ± 0.071	9.98 ± 0.74
40t/ha	17.5 ± 0.85	95.08 ± 3.21	0.75 ± 0.041	15.3 ± 0.50
60t/ha	23.7 ± 1.12	102.6 ± 2.86	0.95 ± 0.088	20.3 ± 0.31
80t/ha	25.2 ± 0.84	144.0 ± 4.58	1.35 ± 0.054	27.9 ± 0.59

• Mean value ± Standard Error

The levels of the various fractions of each metal increased with increasing the sewage sludge application rate, i.e., followed the same pattern of the total content. The data in Table (3) showed that the soluble and exchangeable fractions (F1) represented only a small percentage of the total content, averaged to be 0.62 ± 0.23 , 1.52 ± 0.65 , 1.77 ± 0.67 , $4.13.\pm 1.65$ % for Cu, Zn, Pb, and Cd, respectively. These lower percentages indicate that heavy metals applied to the soil from sewage sludge were strongly sorbed in non-exchangeable form. Aboulroos et al., (1991); Badawy and Helal

(1997) showed that the soluble and exchangeable fractions represented only 1-5% from the total Cd, Co, Ni, Pb, Cu, and Zn content of the soil irrigated with sewage effluent. The carbonate fraction (F2) of heavy metals represented an average of 7.47 ± 0.85 , 10.1 ± 0.66 , 9.1 ± 1.74 , $10.34 \pm 0.81\%$ for Cu, Zn, Pb, and Cd, respectively, The Fe and Mn oxides fraction (F3) of heavy metals represented an average of 6.4 ± 0.95 , 1.6 ± 3.39 , 23.9 ± 4.40 , $13.7 \pm 3.28\%$ from the total content for Cu, Zn, Pb, and Cd, respectively. The organic fraction (F4) of heavy metals represented an average of 35 ± 2.14 , 26 ± 1.20 , 22 ± 1.11 , $30 \pm 1.95\%$ from total content for Cu, Zn, Pb, and Cd, respectively. Also, the residual fraction of these metal approximately represented from 42 to 51% from the total content. At any given time the order of Cu, Zn, Pb, and Cd in the various fractions are as follows: residual> organic> oxides> carbonates> exchangeable. Distributions of different fractions of Cu, Zn, Cd, and



Fig.1. Chenges of soil OM, CaCO₃ and pH as a fanction of time and sewage sludge application rates.

Pb expressed as percentages of the total and in relation with the time factor are given in Figure (2). This indicates that the soluble and exchangeable fractions and organic fraction increased

Fraction*	(Cu	Z	Zn	
	Range Average \pm SD		Range	Average \pm SD	
F1	0.3 – 1.0	0.62 ± 0.23	0.5 - 2.5	1.52 ± 0.65	
F2	6.0 - 8.4	7.42 ± 0.85	9.0 - 11	10.1 ± 0.66	
F3	4.8 - 8.9	6.35 ± 0.55	14 - 23	17.1 ± 3.39	
F4	32 - 38	34.8 ± 2.14	25 - 28	26.4 ± 1.20	
F5	47 - 51	49.2 ± 4.67	50 - 53	51.2 ± 2.75	
	(Cd	P	b	
	Range	Average \pm SD	Range	Average \pm SD	
F1	2.0 - 7.0	4.13 ± 1.65	0.5 - 2.5	1.77 ± 0.67	
F2	7.0 - 12.0	9.10 ± 1.74	9.0 - 11	10.3 ± 0.81	
F3	18 - 28	23.9 ± 4.40	14 - 23	13.7 ± 3.28	
F4	21 - 24	21.9 ± 2.11	25 - 28	29.7 ± 1.95	
F5	40 - 44	41.8 ± 2.33	50 - 53	50.2 ± 1.67	

Table (3): percentage from total content of different fraction of various metals in soil treated with different application rates of sewage sludge.

*F1.= exchangeable; F2 = bound to carbonates; F3 = bound to Fe and Mn oxides; F4 = bound to organic matter; F5= Residual

with time up to 6 months then decreased gradually. The carbonates fraction decreased according to Aboulroos et. al., (1989) due to the release of organic acids from sewage sludge, whereas, the oxides fraction increased with time. This may be due to metal transformation from soluble and exchangeable to oxide form, Aboulroos et al., (1991).

The relative distribution shows that the concentrations of heavy metals decreased in the mobile fractions with time, {(compare 0 time (start) with 12 months (end)}, approximately 29, 17, 51, 43 % of Cu, Zn, Cd, and Pb, respectively. The exchange of Cu, Zn, Cd, and Pb between mobile and inert fractions was estimated by calculating a mobility factor (MF) {the ratio of mobile to inert fractions}. Selected mobility factor for cultivated soil at 0 time (start), 3, 6, 9, and 12 month (at end) are presented in Table (4). This indicate that the value of MF increased with increasing the time, up to 6 month then it decreased with time. It should be mentioned that low values of MF indicate low mobility of the metals. Almas et al. (1999) found that the concentrations of mobile fractions decreased and of inert fractions increased with time in soil treated with organic matter. The MF of Cd is higher than other metals (Cd > Pb > Zn > Cu) this due to the organic complextation of Cd is lower than this metals (Fig. 3).

Time	MF					
(months)	Cu	Zn	Cd	Pb		
0 (start)	0.088	0.127	0.168	0.139		
3	0.099	0.137	0.169	0.141		
6	0.150	0.159	0.220	0.196		
9	0.067	0.108	0.129	0.104		
12	0.048	0.067	0.099	0.064		

Table (4): Mean values of mobility factors (MF) for soil heavy metal contents at different time of starting the experiments with 80 t/ha application rates sewage sludge.



Fig. 2. Relative distrbution of Cu, Zn, Cd, and Pb among the different fractions of the soil treated with 80 t/ha sewage sludge, at icreasing equilibrium time.

DTPA-extractable heavy metals:

Data in Table (5) show a substantial increase in the DTPA extractable metals due to the increase in sewage sludge application rate, followed the same pattern observed with the total metals content. Comparing the values of the highest treatment of sewage sludge with the control, these increased in the order: Cu 17 times < Zn 12 times < Pb 11 time < Cd 6 times. The DTPA- extractable Cu, Zn, Cd, and Pb are negatively correlated with soil pH (r = -0.837**, -0.915**, -0.644**, and -0.884**, respectively), soil CaCO₃ content (r = -0.519**, -0.668**, -0.594*, -0.674**, respectively), and positively correlated with soil organic matter content (0.544**, 0.573**, 0.545**, 0.595**, respectively).

The percentage of the total content of heavy metals extracted by DTPA (Table 6) varied widely from one metal to another, indicating wide variations in their relative availability. These percentages, increased in the order: Zn 4% < Cu 8.5% < Cd 9% < Pb 10% at the highest sewage sludge application rate. Figure (4) shows the continuous increase in soil DTPA extractable Cu, Zn, Cd, and Pb with increasing the sewage sludge application rate. The maximum increase happened at 6 months in relation to the decrease in soil pH and CaCo₃. The availability of these metals to plants decreased after 6 months, indicting transformation of a part of accumulated metal from available to unavailable form with the time. A glance at the data in Table (6) indicated that vegetation increased the percentages of DTPA extractable Cu, Zn, Cd, and Pb comparing with uncultivated soils up to 6 months, after which a reduction of these metals occurred, due to plant uptake In uncultivated soil they decreased, but still more than the cultivated soil.



Table (5): DTPA extractable-metals (mg/kg soil) in soil amended with different sewage sludge application rates at different periods time of time.

Metal (mg/kg soil)		Time (month)									
		0	3	6	9	12	0	3	6	9	12
Treatme	ent			Cu			Zn				
	Control	0.17	0.26	0.40	0.45	0.51	0.48	0.45	0.42	0.30	0.61
	20t/ha	0.20	0.18	4.06	1.32	1.22	2.52	4.20	6.13	4.65	4.07
Cultivated	40t/ha	2.66	4.17	5.61	2.07	1.97	4.52	6.00	8.45	5.98	5.36
	60t/ha	2.65	5.30	6.96	2.50	2.80	5.23	7.60	10.15	6.63	6.17
	80t/ha	3.46	5.44	8.52	3.02	3.83	6.40	8.87	13.82	7.98	7.83
	Control	0.17	0.40	0.78	0.65	0.70	0.48	0.35	0.42	0.48	0.45
Uncultivated	20t/ha	0.20	0.29	1.84	3.00	2.67	2.52	3.70	5.13	5.40	4.75
	40t/ha	2.66	3.10	4.65	3.93	3.12	4.52	5.80	7.45	6.30	5.67
	60t/ha	2.65	3.79	5.13	4.30	3.1	5.23	5.50	8.40	7.80	7.50
	80t/ha	3.46	4.94	6.50	5.94	4.62	6.40	6.15	10.75	9.25	8.25
		Cd					РЬ				
	Control	0.02	0.03	0.03	0.01	0.01	0.23	0.25	0.27	0.25	0.24
	20t/ha	0.04	0.10	0.14	0.10	0.10	0.97	1.40	2.29	1.45	1.25
Cultivated	40t/ha	0.07	0.11	0.20	0.13	0.13	1.25	2.57	3.13	1.98	1.75
	60t/ha	0.80	0.14	0.30	0.16	0.15	2.11	2.93	4.45	2.50	2.47
	80t/ha	0.12	0.20	0.42	0.22	0.22	2.69	3.65	5.54	3.35	3.27
	Control	0.02	0.03	0.04	0.01	0.03	0.26	0.21	0.25	0.27	0.26
Uncultivated	20t/ha	0.05	0.06	0.14	0.12	0.11	0.97	1.29	2.04	1.85	1.39
	40t/ha	0.07	0.09	0.16	0.15	0.14	1.25	1.70	2.50	2.40	2.10
	60t/ha	0.08	0.11	0.20	0.18	0.17	2.11	2.53	3.13	3.47	3.00
	80t/ha	0.11	0.15	0.29	0.24	0.22	2.67	3.01	4.70	4.20	4.00

Time	Cu		Zn			Cd	Pb	
(month)								
	Cultivated	Uncultivated	Cultivated	Uncultivated	Cultivated	Uncultivated	Cultivated	Uncultivated
0(start)	9.35	8.05	4.05	4.27	9.05	8.17	9.78	10.27
3	14.7	13.54	5.61	4.16	15.5	11.11	13.27	11.58
6	23.06	17.50	8.75	7.17	32.62	21.42	20.45	18.08
9	9.17	16.05	5.05	6.17	17.05	17.89	12.18	16.15
12	10.36	12.49	4.97	5.50	17.00	16.35	11.89	15.38

Table (6): Changes in soil DTPA-extractable metal as a percentage from total contents.

Heavy metals in plant:

Data in Table (7) show that the concentration of Cu, Zn, Cd, and Pb in leaves and fruits increased with increasing sewage sludge application rate to the soil. The results indicate that the concentration of heavy metals in the leaves are higher than the fruits. The absolute values of these metals in the leaves are different from metal to another. This depends on the sewage sludge content and the availability of these metals in the soil which are controlled by different soil parameters (CaCO₃, pH...etc.). The values of these metals followed the order Zn > Cu > Pb > Cd. The increase in metals content at the highest sludge application rate compared with the control was 6 for Cu, 2.5 for Zn, 14 for Cd, and 20 fold for Pb; for leaves and 5 for Cu, 2 for Zn, 31 for Cd, and 25 fold for Pb for fruits, respectively. The concentration of Cd in tomato leaves and fruits are higher than normal. The intermediate concentration of Cd reported by Adriano (1986) was <0.30 ppm for leaves and 0.04-0.06 ppm for fruits. However, the concentration of Pb is within the normal range compared with tomato leaves grown in uncontaminated soil (6.3 ppm for leaves and 1.19 ppm for fruits) as reported by Chapman (1975). A highly significant correlation coefficient was found between both leaves and fruits, and soils DTPA-extractable content of Cu, Zn, Cd, and Pb (0.964** and 0.912**; 0.914** and 0.945**; 0.867** and 0.794**; and 0.887** and 0.909**, respectively).

	Cu		Zn		Cd		Pb			
treatment	leaves	fruits	leaves	fruits	leaves	fruits	leaves	fruits		
control	3.8	2.00	28.9	22.5	0.09	0.04	0.14	0.05		
20 t/ha	12.8	6.52	45.8	30.5	0.54	0.39	1.42	0.56		
40 t/ha	15.6	8.34	69.5	35.8	1.10	0.68	1.85	0.69		
60 t/ha	18.9	9.20	75.9	38.8	1.34	0.85	2.54	0.84		
80 t/ha	22.5	10.58	86.5	42.5	1.75	1.25	3.25	1.25		
L.S.D.0.05	2.92	1.55	4.56	2.58	0.21	0.14	0.27	0.08		

Table (7): Heavy metal concentrations (mg/kg dry matter) in different tissues of tomato plants grown on soil treated with different rates of sewage sludge.



Fig.4. DTPA extractable-metals as a fancution of time and sewage sludge application rates.

Acknowledgment

The authors wish to thank the International Atomic Energy Agency for the financial support provided through the TC project No EGY/8/013

References

- Aboulroos, S.A., Holah, Sh.Sh, and Badawy, S.H. (1989). Influence of prolonged use of sewage effluent in irrigation on heavy metal accumulation in soils and plants. Z. Pflanzenernahr. Bodenk. 152: 51-55.
- Aboulroos, S.A., Holah Sh.Sh., El-Kherbawy, M.I, and Badawy, S.H. (1991). Fractionation of some heavy metals in soils irrigated with sewage effluent for different years. Egypt. J. Soil Sci. 31: 43-55.
- Aboulroos, S.A., Holah Sh.Sh., and Badawy, S.H. (1996). Background levels of some heavy metals in soil and corn in Egypt. J. Soil Sci. 36: 83-97.
- Adriano, D.C. (1986). Trace elements in the terrestrial environment. Springer-verlage, New York.
- Almas, A., Singh, B.R., and Salbu, B. (1999). Mobility of cadmium-109 and zinc 65 in soil influenced by equilibrium time, temperature, and organic matter. J. Environ. Qual. 28: 1724-1750
- Badawy, S.H., and Helal, M.I.D. (1997). Impacts of heavy metals of sewage effluent on soils and plants of Helwan area. J.Agric.Sci. Mansoura Univ. 22:4737-4754.
- Chapman, H.D. (1975). Diagnostic criteria for plant and soils. Eurasia Publ. House (p) LTD. Ram Nagar, New Delhi-110055
- Christensen, T. H., and Huang, P.M. (1999). Solid phase cadmium and the reactions of aqueous cadmium with soil surfaces. In M.J. McLaughlin and B.R. Singh (ed.) cadmium in soil and plants. Kluwer Academic Publ. Dordecht, Netherlands.
- Dewis J. and Freites F. (1970). Physical and chemical methods of soil & water analysis. FAO. Rome. Soil Bulletin No. 10.
- El-Motuim, R.A, and Badawy, S.H. (2000) Effect of irrigation using sewage water on the distribution of some heavy metals in bulk and Rhizosphere soils and different plant species: Cabbage plants (Brassica Oleracea L.); Orange tree (Citrus Sinenss L.). Egypt. J. Soil Sci. (under press).
- El-Sokkary, I.H. (1979). The chemical fractionation of soil zinc and its specific and total absorption by Egyptian alluvial soil. Plant and Soil. 53: 117-129.
- He, Q.B., and Singh, B. R. (1993). Effect of organic matter on the distribution, extractability and uptake of cadmium in soil. J. Soil Sci. 44: 641-650.
- Jackson, M.L. (1967). Soil chemical analysis. Prentice-Hall India Part. Ltd., New Delhi, India.
- Kennedy, V.H., A.L. Sanchez, D.H. Oughton, and Rowland, A.P. (1997). Use of single and sequential chemical extraction to assess radionuclide and heavy metal availability from soils for root uptake. Analyst. 122: R89-R100.
- Lindsay, W.L., and Norvell, W.A. (1978). Development of DTPA soil test for zinc, iron, manganese, and copper. Soil Sci. Soc. Am. J. 42: 421-428.
- McBride, M.B. (1995). Toxic metal accumulation from agricultural use of sludge: are USEPA regulations protective. J. Environ. Qual. 24: 5-18
- Narwal, R.P., and Singh , B.R. (1998). Effect of organic materials on partitioning, extractability and plant uptake of metals in an alum shale soil. Water Air Soil Pollut. 103: 405-421.
- Pickering, W.F. (1998). Metal ion speciation-soils and sediments. Ore Geol. Rev. 1: 83-146.

- Reisenauer H.M. (1978). Soil and Plant-tissue testing in California. Division of Agriculture Sciences, University of California.
- Salbu, B., T. Krekling, and D.H. Oughton (1998). Characterisation of radioactive particles in the environment. Analyst. 123: 843-849.
- Singh, B.R., R.P. Narwal, and Almas, A.R. (1995). Crop uptake and extractability of cadmium in soils naturally high in metals art different pH levels. Commun. Soil Sci. Plant Anal. 26: 2123-2142.
- Singh, B.R., R.P. Narwal, and Almas, A.R. (1998). Residual effect of organic mater on cadmium uptake by plants and its distribution in soils. In R. Rost (ed.) Third international conference on Bio-geochemistry of Trace Elements. Contaminated Soils, Paris. CD-Room D:\ communic\019pdf.
- Sposito, G., Levesque, C.S., Leclaire, J.P., and Chang, A.C. (1983). Trace metal chemistry in aridzone field soils amended with sewage sludge. III. Effect of time on the extraction of trace metals. Soil Sci. Soc. Am. J. 47: 898-902.
- Tessier, A., Campbell, P. G. C., and Bisson, M. (1979). Sequential extraction procedure for the speciation of particulate trace metals. Anal. Chem. 51: 844 -850.